

TYLENOL (ACETAMINOPHEN) DOSAGE CHART

Note: Tylenol can be given every 4 hours. DO NOT give to children under 2 months of age.

WEIGHT (lbs)	DOSE (mgs)	INFANT ORAL SUSPENSION (160 mg/5 ml)	CHILD ORAL SUSPENSION (160 mg/5 ml)	CHEWABLE (80 mg tab)	JR. STRENGTH (160 mg tab)	ADULT TAB (325 mg)
6 - 11	40	1.25 ml=1/4 tsp	-	-	-	-
12-17	80	2.5 ml=1/2 tsp	2.5 ml=1/2 tsp	-	-	-
18-22	120	3.75 ml=3/4 tsp	3.75 ml=3/4 tsp	1 1/2 tabs	-	-
23-35	160	5 ml=1 tsp	5 ml=1 tsp	2 tabs	1 tab	-
36-47	240		7.5 ml = 1 1/2 tsp	3 tabs	1 1/2 tab	-
48-59	320		10 ml = 2 tsp	4 tabs	2 tabs	1 tab
60-71	400		12.5 ml = 2 1/2 tsp	5 tabs	2 1/2 tabs	1 tab
72-95	480		15 ml = 3 tsp	6 tabs	3 tabs	1 1/2 tab
96+	640		20 ml = 4 tsp	8 tabs	4 tabs	2 tabs

MOTRIN & ADVIL (IBUPROFEN) DOSAGE CHART

Note: Ibuprofen can be given every 6 hours. DO NOT give to children under 6 months of age.

WEIGHT (lbs)	DOSE (mgs)	INFANT ORAL DROPS (50 mg/1.25 ml)	CHILD ORAL SUSPENSION (100 mg/5 ml)	CHEWABLE (100 mg tab)	JR. STRENGTH (100 mg tab)	ADULT TAB (200 mg)
6 - 11	-	-	-	-	-	-
12-17	50	1.25 ml=1 dropper	2.5 ml=1/2 tsp	-	-	-
18-22	75	1.875 ml=1 1/2 dropper	3.75 ml=3/4 tsp	1 tab	1 tab	-
23-35	100	2.5 ml=2 dropper	5 ml=1 tsp	1 tab	1 tab	-
36-47	150		7.5 ml = 1 1/2 tsp	1 1/2 tabs	1 1/2 tab	-
48-59	200		10 ml = 2 tsp	2 tabs	2 tabs	1 tab
60-71	250		12.5 ml = 2 1/2 tsp	2 1/2 tabs	2 1/2 tabs	1 tab
72-95	300		15 ml = 3 tsp	3 tabs	3 tabs	1 1/2 tab
96+	400		20 ml = 4 tsp	4 tabs	4 tabs	2 tabs

BENADRYL (DIPHENHYDRAMINE) DOSAGE CHART

Note: Benadryl can be given every 6 hours. Consult your physician for children under 12 months of age.

WEIGHT (lbs)	AGE (Years)	SUSPENSION (12.5 mg/5 ml)	CHEWABLE (12.5 mg tab)	FASTMELTS (12.5 mg tab)	TAB/CAP (25 mg)
12-19	1-1 1/2	2.5 ml=1/2 tsp	-	-	-
20-26	1 1/2-2	3.75 ml=3/4 tsp	-	-	-
27-39	2-4	5 ml=1 tsp	1 tab	1 tab	-
40-52	5-6	7.5 ml=1 1/2 tsp	1 tab	1 tab	-
53-67	7-8	10 ml=2 tsp	2 tabs	2 tabs	1 tab/cap
68-79	9-10	10-12.5 ml=2-2 1/2 tsp	2 tabs	2 tabs	1 tab/cap
80-95	11-12	10-15 ml= 2-3 tsp	2-3 tabs	2-3 tabs	1 tab/cap
96+	12+	10-20 ml=2-4 tsp	2-4 tabs	2-4 tabs	1-2 tabs/caps

Caution! Benadryl can cause significant sleepiness or unexpected hyperactivity in some children. Dosing by weight is most accurate but consult your physician for any questions.